



Fact Sheet

South Carolina Department of Health and Environmental Control • www.scdhec.gov

Group A Streptococcal

What is group A strep (GAS) and invasive group A strep?	Group A Streptococcal (GAS) bacteria cause common infections such as strep throat and impetigo. At any time up to 10% of school children may carry the bacteria and have no symptoms. The CDC estimates that there are several million cases of strep throat and impetigo, and about 10 thousand invasive GAS cases each year. Most Group A Streptococcal infections are not serious or invasive and are not reportable to the SC Department of Health & Environmental Control. Invasive Group A Strep infections are rare complications of GAS. These more severe infections include: sepsis (an invasion of bacteria in the bloodstream); pneumonia; necrotizing fasciitis, an infection that affects the deeper layers of skin and underlying tissues; and, rarely, Streptococcal Toxic Shock Syndrome, which starts at the site of a wound.
What are the symptoms of invasive Group A Strep?	Symptoms will vary depending on the strain of GAS and where the bacteria has entered the body. <u>Sepsis</u> – can produce high fever, shock, inflammation of internal organs, or abscesses. <u>Necrotizing Fasciitis</u> – the infected skin is painful, hot and swollen and the patient usually has severe body pain, high fever, and rapid pulse. Sepsis may develop. This strain of GAS destroys skin & deep tissue and is sometimes referred to by the media as “flesh-eating bacteria”. <u>Toxic Shock Syndrome</u> – begins with flu like symptoms such as fever, chills and muscle pain, rash, low blood pressure (shock), kidney malfunction and blood and liver abnormalities. (Note: this is not the Toxic Shock Syndrome usually associated with tampon use and Staph bacteria.)
How is it treated?	People with sore throats, impetigo, or infected wounds should see their doctor for medical care and possible Strep testing and antibiotics. Symptoms of invasive forms of Group A Strep are very serious and require immediate medical attention. Diagnosis can sometimes be difficult to make early in the course of infection. Treatment includes antibiotics, intensive medical care, and in some cases, surgery.
How do people catch this disease?	Group A Strep can be spread through direct, close personal contact with the oral secretions or infected tissue fluids of someone with GAS infection. Most people who are exposed to Group A Strep do not get infected or become ill. Only close personal contacts (ie, household members) who are exposed to invasive GAS strain are at minimal risk of being infected.
What can be done to prevent this disease?	Careful and regular handwashing is the best prevention, especially before eating, after coughing or sneezing or using the bathroom, and before preparing foods. Cuts and abrasions should be kept clean and covered. Anyone who has been diagnosed with non-invasive group A strep should stay home from work or school until they have been on antibiotics for at least 24 hours. Avoid close personal contact with anyone who has strep throat, wound infection or impetigo or GAS infection until 1-2 days after antibiotics are begun.
Is the public at risk?	There is no increased risk to the general public. Even close personal contacts (household, personal care workers) are at only minimal increased risk.

SOUTH CAROLINA DEPARTMENT OF HEALTH AND ENVIRONMENTAL CONTROL

Region 7

Serving Berkeley, Charleston and Dorchester Counties

Public Health Office • 4050 Bridge View Drive • Suite 600 • N. Charleston, SC 29405 • Phone: (843) 746-3800 • www.scdhec.gov